HWB10

Oxfordshire Health & Wellbeing Board

18th June 2020

Oxfordshire Health & Wellbeing Joint Strategic Needs Assessment 2020

Introduction

The Joint Strategic Needs Assessment (JSNA) was introduced to create stronger partnerships between communities, local government and the NHS, providing a firm foundation for commissioning that improves health and social care provision and reduce health inequalities. It helps commissioners shape services to address local needs, and a number of supportive resources are used to help produce the JSNA.

Since 2013 the Health & Wellbeing board has responsibility of overseeing the production of the JSNA. The JSNA is the means by which local leaders work together to understand and agree the needs of all local people, with the joint health and wellbeing strategy setting the priorities for collective action. Taken together they are the pillars of local decision making, focussing on the priorities for action and providing the evidence base for decisions about local services.

The JSNA in 2020

The JSNA is a draft working document published in draft form on the Oxfordshire Insight webpage through the following link

https://insight.oxfordshire.gov.uk/cms/system/files/documents/2020_JSNA_DRAFT.pdf

Building on previous years, the team have developed the format of the document this year which makes it more user friendly and easier to search.

The report continues to be a mix of national statistics, local statistics and links to research carried out by organisations in Oxfordshire.

The report this year has more information on inequalities and more use of maps.

The datasets were produced pre the coronavirus pandemic. Due to the pandemic some national data collection had been paused so there will be delays in updating some of the national datasets.

Key findings of the Report

- Oxfordshire overall is relatively healthy, but inequalities remain;
 - Oxfordshire's more deprived wards experience significantly higher mortality rates than England.
 - The gap in early years development between lower income pupils in Oxfordshire has increased for the second year in a row.
- Just over 2,000 deaths in Oxfordshire (over three years 2016-18) were considered preventable including 1,054 cancer deaths and 558 cardiovascular.

HWB10

- Almost two thirds of adults are classified as overweight or obese. Prevalence is higher in males, older people, some ethnic groups and in more deprived areas.
- Oxfordshire's first health new towns Bicester and Barton have highlighted strong partnership working and people actively managing their health.
- There was an increase in the number of police recorded victims of domestic abuse in Oxfordshire and a significant increase in domestic abuse crimes involving children.
- Mental health rates of diagnosis and referrals are continuing to increase.

A range of facts and figures from the JSNA are shown in the diagram below.

Recommendation

The board are asked to sign off the draft JSNA report for 2020.

Oxfordshire JSNA, health and wellbeing facts and figures 2020





	. ()) _	_	i,	data.	*	*	_	_		£	
PRE-BIRTH	7.5		SIRTH	7,402	687,500	82.5%		%	EARLY	74.0%	10%	
	Moth	ers		Babies (Oxfordshire	Mothers st	art Babies br	eastfed	YEARS G	ood level of	Children live in	
-00	smoking:	at birth		born	population	breastfeedin birth	gat at 6-8 w	eeks		opment by end undation stage	low income families	
4) (M	<u>₹</u>		1					•		Ζ,
779	9 4	6.6%	126	6,464	164		ONDARY	65%	1,987	1,34		
Looked				ung people	Not in educ	cation,				ren Reception d		
After Childre	_	and Maths	1	ith learning	employme trainin			reading, writin	g, overweight obese	or overweigh obese		
Unildre				difficulties		•		maths	400	-		
	1		78		4			Ĥ	£ŧ	£Ħ		
121	. 5	58	165	1,147	7,3	54 W	ORKING AGE	14,985	£30,428	£37,250	6,100	
Under	18 Young	people Un	der 18s hospital	15-24s	Apprent	iceship	AGE	everage	average	average	Adults are	
concepti			dmissions for	diagnosed wi		pants	he	use price	female salary	male salary	unemployed	
_			cohol (3 years)	Chlamydia	400		ملم		_		A	
Æ	•	4	- 2	7		್)	•	*	A	
273	3	1,701	17,	857	67,577	72.5%	10,6	31	59%	54,804	304	
Killed or se	riously Peo	ple supporte	d with Adult	s with	Adults	Adults an	_		ults overweight		omeless households	
injured on	roads le	arning disabi			gnosed with		y obstructive p disea		or obese	smoke	in priority need	
					depression	active				43.0	-	
T	3		2		<mark></mark>	* FLEST	.	<u> </u>	7	A.	Ø	
1,4	27	155	4,493	89	,800	48	3,001	11,7	779	23,438	31,146	
Alcohol	nospital	Deaths from	Deaths fr	om Adults		Deaths from dr	ug New Sexua			useholds in	Adults with	
admissio	_	uicide (3 year:	s) cancer (3 y	,	illness or	misuse (3 year			ic abuse fu	el poverty	diabetes	
40-			400		ability		Infection:		400	-	Alla	
	\$	+#	+1		•	<i>\$</i>	%		##	ΨŤ	11	
END OF LIFE	51%	84.7	81.6	5,8	52 1	13,134	2,750	OLDER PEOPLE	71.7	68.3	61,100	
	ied in usual	Average	Average		with Per	ople had a	Hospital		Average female	Average ma		
	place of	female life			entia	stroke a	lmissions due		healthy life	healthy lif		
	residence	expectancy	expectan	cy			to falls		expectancy	expectano	У	